

Screening for STIs, HIV, and Hepatitis B/C during Pregnancy in NYS

Infection	1st Prenatal Visit	3rd Trimester	Delivery
Syphilis [^]	Everyone	Everyone	Everyone
Gonorrhea*	If <25, or ≥25 at risk	If at risk	N/A
Chlamydia*	If <25, or ≥25 at risk	If <25, or ≥25 at risk	N/A
HIV	Everyone	Everyone	If not tested during this pregnancy or at risk
Hepatitis B [^]	Everyone	N/A	If not tested during this pregnancy or at risk
Hepatitis C [^]	Everyone	If not tested during this pregnancy or at risk	If not tested during this pregnancy or at risk

Screening Test
Syphilis Serology
GC NAAT
CT NAAT
HIV-1/2 Ab/Ag
HBsAg
HCV Ab with reflex to HCV RNA

[^]Testing mandated by NYS Public Health Law.

*Repeat screen 3 months after treatment of documented gonorrhea or chlamydia infection during pregnancy.

Risk factors for STI/HIV and hepatitis B and C may include: History or current diagnosis of an STI; new partner(s); pregnant person or partner with multiple partners; sex partner with an STI; condomless sex not in a tested negative mutually monogamous relationship; transactional sex; history of incarceration; pregnant person or partner with injection drug use; high incidence/prevalence setting.

Compiled from NYS, USPSTF, and CDC screening guidelines and NYS public health laws and regulations.

Clinical Questions: CEI Line 866-637-2342

